

GEOMETRY OF INFINITE DIMENSIONAL CARTAN DEVELOPMENTS

JOHANNA MICHOR AND PETER W. MICHOR

ABSTRACT. The Cartan development takes a Lie algebra valued 1-form satisfying the Maurer-Cartan equation on a simply connected manifold M to a smooth mapping from M into the Lie group. In this paper this is generalized to infinite dimensional M for infinite dimensional regular Lie groups. The Cartan development is viewed as a generalization of the evolution map of a regular Lie group. The tangent mapping of a Cartan development is identified as another Cartan development.

1. INTRODUCTION

A regular Lie group G is one where where one can integrate smooth curves in the Lie algebra \mathfrak{g} to smooth curves in the Lie group in a smooth way: The evolution mapping $\text{Evol} : C^\infty(\mathbb{R}, \mathfrak{g}) \rightarrow C^\infty(\mathbb{R}, G)$ exists and is smooth; see 2.4 below. This notion is relevant for infinite dimensional Lie groups where existence results for ODEs are not available in general. A stronger version of this notion is due to Omori et al. [20, 21, 22, 23, 24, 25] and was weakened to this version for Fréchet Lie groups by Milnor [19]. It was generalized to Lie groups modeled on convenient vector spaces (i.e., locally convex vector spaces where each Mackey Cauchy sequence converges) in [14], see also [13]. Up to now, no example of a non-regular Lie group modelled on convenient vector spaces is known. In section 2 we review regular Lie groups in order to fix notation and to make clear the analogy to the Cartan-development.

One can extend the notion of regularity to other classes of curves for certain Lie groups (e.g., modeled on Banach spaces) and ask for the existence of Evol on $C^k(\mathbb{R}, \mathfrak{g})$ for $k = 0, 1, 2, \dots$ or even $L^1(\mathbb{R}, \mathfrak{g})$. Results in these directions are due to Glöckner in [6, 7] and Hanusch in [11, 10, 9].

In this paper we extend the domain of the evolution mapping: In order to end up in $C^\infty(M, G)$ for an infinite dimensional manifold M we consider the spaces $\Omega_{\text{flat}}^1(M, \mathfrak{g})$ of flat \mathfrak{g} -valued differential 1-forms on M in 3.1 and show that the evolution map exists and is smooth for simply connected pointed M , and we give an explicit description of its tangent mapping. This is an extension to infinite dimensions of the Cartan development.

2020 *Mathematics Subject Classification.* Primary 58B25.

Key words and phrases. Cartan development, regular infinite dimensional Lie group.

Johanna Michor was supported by FWF-Project P31651.

The results of this paper could be applied in situations where a diffeomorphism group acts on a space of shapes; the diffeomorphism mapping a template shape to any shape could be interpreted as a Cartan-development. See also [4].

Infinite dimensional manifolds Lie groups are always modeled on convenient vector spaces as treated in [13], see also [18] or [29]

This paper was inspired by the paper [5] on integrable surfaces which was pointed out to the authors by Oleksandr Sakhnovich. In [5] the term ‘integrable’ just means that the 1-forms considered there satisfy the Maurer-Cartan formula, and it has nothing to do with complete integrability: There is no Hamiltonian aspect to this, as was pointed out to us by Boris Khesin.

2. REVIEW OF REGULAR LIE GROUPS

2.1. Notation on Lie groups. Let G be a Lie group which may be infinite dimensional with Lie algebra \mathfrak{g} . Later we will only consider regular Lie groups, see 2.4. Let $\mu : G \times G \rightarrow G$ be the multiplication, let μ_x be left translation and μ^y be right translation, given by $\mu_x(y) = \mu^y(x) = xy = \mu(x, y)$. We denote inversion by $\nu : G \rightarrow G$, $\nu(x) = x^{-1}$. The tangent mapping $T_{(a,b)}\mu : T_aG \times T_bG \rightarrow T_{ab}G$ is given by

$$T_{(a,b)}\mu.(X_a, Y_b) = T_a(\mu^b).X_a + T_b(\mu_a).Y_b$$

and $T_a\nu : T_aG \rightarrow T_{a^{-1}}G$ is given by

$$T_a\nu = -T_e(\mu^{a^{-1}}).T_a(\mu_{a^{-1}}) = -T_e(\mu_{a^{-1}}).T_a(\mu^{a^{-1}}).$$

Let $L, R : \mathfrak{g} \rightarrow \mathfrak{X}(G)$ be the left and right invariant vector field mappings, given by $L_X(g) = T_e(\mu_g).X$ and $R_X = T_e(\mu^g).X$, respectively. They are related by $L_X(g) = R_{\text{Ad}(g)X}(g)$. Their flows are given by

$$\text{Fl}_t^{LX}(g) = g \cdot \exp(tX) = \mu^{\exp(tX)}(g), \quad \text{Fl}_t^{RX}(g) = \exp(tX).g = \mu_{\exp(tX)}(g).$$

We also need the right Maurer-Cartan form $\kappa = \kappa^r \in \Omega^1(G, \mathfrak{g})$, given by $\kappa_x(\xi) := T_x(\mu^{x^{-1}}) \cdot \xi$. It satisfies the left Maurer-Cartan equation $d\kappa - \frac{1}{2}[\kappa, \kappa]_\wedge = 0$ which follows by evaluating $d\kappa^r$ on right invariant vector fields R_X, R_Y for $X, Y \in \mathfrak{g}$

$$\begin{aligned} (d\kappa^r)(R_X, R_Y) &= R_X(\kappa^r(R_Y)) - R_Y(\kappa^r(R_X)) - \kappa^r([R_X, R_Y]) \\ &= R_X(Y) - R_Y(X) + [X, Y] = 0 - 0 + [\kappa^r(R_X), \kappa^r(R_Y)], \end{aligned}$$

where $[\cdot, \cdot]_\wedge$ denotes the wedge product of \mathfrak{g} -valued forms on G induced by the Lie bracket. Note that $\frac{1}{2}[\kappa, \kappa]_\wedge(\xi, \eta) = [\kappa(\xi), \kappa(\eta)]$.

Similarly the *left Maurer-Cartan form* $\kappa^l \in \Omega^1(G, \mathfrak{g})$ is given by $\kappa_g^l = T_g(\mu_{g^{-1}}) : T_gG \rightarrow \mathfrak{g}$ and it satisfies the right Maurer-Cartan equation $d\kappa^l + \frac{1}{2}[\kappa^l, \kappa^l]_\wedge = 0$. We have also $(\nu^*\kappa^r)_g = -\kappa_g^l = -\text{Ad}(g^{-1})\kappa_g^r$.

The (exterior) derivative of the function $\text{Ad} : G \rightarrow GL(\mathfrak{g})$ can be expressed by

$$d\text{Ad} = \text{Ad} \cdot (\text{ad} \circ \kappa^l) = (\text{ad} \circ \kappa^r) \cdot \text{Ad},$$

since, for $c : \mathbb{R} \rightarrow G$ smooth with $c(0) = e$ and $c'(0) = X$ like $\exp(tX)$ if \exp exists,

$$d\text{Ad}(T\mu_g.X) = \partial_t|_0 \text{Ad}(g.c(t)) = \text{Ad}(g)\partial_t|_0 \text{Ad}(c(t)) = \text{Ad}(g) \cdot \text{ad}(\kappa^l(T\mu_g.X)).$$

Since we shall need it we also note

$$d(\text{Ad} \circ \nu) = -(\text{ad} \circ \kappa^l)(\text{Ad} \circ \nu) = -(\text{Ad} \circ \nu)(\text{ad} \circ \kappa^r).$$

2.2. The right and left logarithmic derivatives. Let M be a manifold and let $f : M \rightarrow G$ be a smooth mapping into a Lie group G with Lie algebra \mathfrak{g} . We define the mapping $\delta^r f : TM \rightarrow \mathfrak{g}$ by the formula

$$\begin{aligned} \delta^r f(\xi_x) &:= T_{f(x)}(\mu^{f(x)^{-1}}).T_x f.\xi_x = \kappa_{f(x)}^r(T_x f.\xi_x) \\ &= (f^* \kappa^r)(\xi_x) \text{ for } \xi_x \in T_x M. \end{aligned}$$

Then $\delta^r f$ is a \mathfrak{g} -valued 1-form on M , $\delta^r f \in \Omega^1(M; \mathfrak{g})$. We call $\delta^r f$ the *right logarithmic derivative* of f , since for $f : \mathbb{R} \rightarrow (\mathbb{R}^+, \cdot)$ we have $\delta^r f(x).1 = \frac{f'(x)}{f(x)} = (\log \circ f)'(x)$.

Similarly the *left logarithmic derivative* $\delta^l f \in \Omega^1(M, \mathfrak{g})$ of a smooth mapping $f : M \rightarrow G$ is given by

$$\delta^l f.\xi_x = T_{f(x)}(\mu_{f(x)^{-1}}).T_x f.\xi_x = (f^* \kappa^l)(\xi_x)$$

2.3. Theorem. *Let $f, g : M \rightarrow G$ be smooth. Then the Leibniz rule holds:*

$$\delta^r(f.g)(x) = \delta^r f(x) + \text{Ad}(f(x)).\delta^r g(x).$$

Moreover, the differential form $\delta^r f \in \Omega^1(M; \mathfrak{g})$ satisfies the ‘left Maurer-Cartan equation’ (left because it stems from the left action of G on itself)

$$\begin{aligned} d\delta^r f(\xi, \eta) - [\delta^r f(\xi), \delta^r f(\eta)]_{\mathfrak{g}} &= 0, \\ \text{or } d\delta^r f - \frac{1}{2}[\delta^r f, \delta^r f]_{\mathfrak{g}} &= 0, \end{aligned}$$

where $\xi, \eta \in T_x M$, and where for $\varphi \in \Omega^p(M; \mathfrak{g}), \psi \in \Omega^q(M; \mathfrak{g})$ one puts

$$[\varphi, \psi]_{\mathfrak{g}}^{\mathfrak{g}}(\xi_1, \dots, \xi_{p+q}) := \frac{1}{p!q!} \sum_{\sigma} \text{sign}(\sigma) [\varphi(\xi_{\sigma_1}, \dots), \psi(\xi_{\sigma_{p+1}}, \dots)]_{\mathfrak{g}}^{\mathfrak{g}}.$$

If $h : N \rightarrow M$ is a smooth mapping, then $\delta^r(f \circ h) = h^*(\delta^r f)$. If $\varphi : G \rightarrow H$ is a smooth homomorphism of groups, then $\delta_H^r(\varphi \circ f) = \varphi' \circ \delta_G^r(f)$ where $\varphi' = T_e \varphi : \mathfrak{g} \rightarrow \mathfrak{h}$.

The left logarithmic derivative the corresponding Leibniz rule is given by

$$\delta^l(fg)(x) = \delta^l g(x) + \text{Ad}(g(x)^{-1})\delta^l f(x).$$

It satisfies the ‘right Maurer Cartan equation’:

$$d\delta^l f + \frac{1}{2}[\delta^l f, \delta^l f]_{\mathfrak{g}} = 0.$$

For ‘regular Lie groups’ a converse to this statement holds; see [13], 40.2. We shall review this in 2.3 and 2.4 below. This result has a geometric interpretation in principal bundle geometry for the trivial principal bundle $\text{pr}_1 : M \times G \rightarrow M$ with right principal action. Then the submanifolds $\{(x, f(x).g) : x \in M\}$ for $g \in G$ form a foliation of $M \times G$ whose tangent distribution is complementary to the vertical bundle $M \times TG \subseteq T(M \times G)$ and is invariant under the principal right G -action. So it is the horizontal distribution of the principal connection $\omega^l \in \Omega^1(M \times G, \mathfrak{g})$ which is given by $\omega^l(\xi_x, \eta_a) := -\delta^l(f.a)(\xi_x) + \kappa^l(\eta_a)$. Thus this principal connection

has vanishing curvature which translates into the result for the right logarithmic derivative.

Proof. For the Leibniz rule we compute for $\xi_x \in T_x M$, using 2.1,

$$\begin{aligned} \delta^r(f \cdot g)(\xi_x) &= T_{f(x).g(x)}(\mu^{(f(x).g(x))^{-1}}).T(\mu \circ (f, g)).\xi_x \\ &= T_{f(x).g(x)}(\mu^{(f(x).g(x))^{-1}}).T\mu.(T_x f.\xi_x, T_x g.\xi_x) \\ &= T(\mu^{f(x)^{-1}}).T(\mu^{g(x)^{-1}}).(T(\mu_{f(x)}).T_x g.\xi_x + T(\mu^{g(x)}).T_x f.\xi_x) \\ &= T(\mu^{f(x)^{-1}}).T(\mu_{f(x)}).T(\mu^{g(x)^{-1}}).T_x g.\xi_x + T(\mu^{f(x)^{-1}}).T_x f.\xi_x \\ &= \text{Ad}(f(x)).\delta^r g(\xi_x) + \delta^r f(\xi_x). \end{aligned}$$

The Maurer-Cartan equation for $\delta^r f$ is the pullback of the one for κ^r :

$$\begin{aligned} d(\delta^r f) &= d(f^* \kappa^r) = f^*(d\kappa^r) = f^*(\tfrac{1}{2}[\kappa^r, \kappa^r]_{\mathfrak{g}}^\wedge) = \tfrac{1}{2}[f^* \kappa^r, f^* \kappa^r]_{\mathfrak{g}}^\wedge \\ &= \tfrac{1}{2}[\delta^r f, \delta^r f]_{\mathfrak{g}}^\wedge \end{aligned}$$

Finally, $\delta^r(f \circ \varphi) = d(f \circ \varphi)^* \kappa^r = d\varphi^* f^* \kappa^r = \varphi^* \delta^r(f)$. For the left logarithmic derivative the proof is analogous. \square

2.4. The 1-dimensional evolution operator. Let G be a possibly infinite dimensional Lie group with Lie algebra \mathfrak{g} . For a closed interval $I \subset \mathbb{R}$ and for $X \in C^\infty(I, \mathfrak{g})$ we consider the ordinary differential equation

$$(1) \quad \begin{cases} g(t_0) = e \\ \partial_t g(t) = T_e(\mu^{g(t)})X(t) = R_{X(t)}(g(t)) \quad \text{or} \quad \kappa^r(\partial_t g(t)) = X(t), \end{cases}$$

for local smooth curves g in G , where $t_0 \in I$. Then the following results hold; see [13], 40.2:

- (2) *Local solution curves g of the differential equation (1) are unique.*
- (3) *If for fixed X the differential equation (1) has a local solution near each $t_0 \in I$, then it has also a global solution $g \in C^\infty(I, G)$.*
- (4) *If for all $X \in C^\infty(I, \mathfrak{g})$ the differential equation (1) has a local solution near one fixed $t_0 \in I$, then it has also a global solution $g \in C^\infty(I, G)$ for each X . Moreover, if the local solutions near t_0 depend smoothly on the vector fields X then so does the global solution.*
- (5) *If (4) holds, then the curve $t \mapsto g(t)^{-1}$ is the unique local smooth curve h in G which satisfies*

$$\begin{cases} h(t_0) = e \\ \partial_t h(t) = T_e(\mu_{h(t)})(-X(t)) = L_{-X(t)}(h(t)) \\ \quad \text{or} \quad \kappa^l(\partial_t h(t)) = -X(t). \end{cases}$$

2.5. Regular Lie groups. If for each $X \in C^\infty(\mathbb{R}, \mathfrak{g})$ there exists $g \in C^\infty(\mathbb{R}, G)$ satisfying

$$(1) \quad \begin{cases} g(0) = e, \\ \partial_t g(t) = T_e(\mu^{g(t)})X(t) = R_{X(t)}(g(t)) \\ \quad \text{or} \quad \kappa^r(\partial_t g(t)) = \delta^r g(\partial_t) = X(t), \end{cases}$$

then we write

$$\begin{aligned} \text{evol}_G^r(X) &= \text{evol}_G(X) := g(1), \\ \text{Evol}_G^r(X)(t) &:= \text{evol}_G(s \mapsto tX(ts)) = g(t), \end{aligned}$$

and call it the *right evolution* of the curve X in G . By 2.4 the solution of the differential equation (1) is unique, and for global existence it is sufficient that it has a local solution somewhere. Then

$$\text{Evol}_G^r : C^\infty(\mathbb{R}, \mathfrak{g}) \rightarrow \{g \in C^\infty(\mathbb{R}, G) : g(0) = e\}$$

is bijective with inverse the right logarithmic derivative δ^r .

The Lie group G is called a *regular Lie group* if $\text{evol}^r : C^\infty(\mathbb{R}, \mathfrak{g}) \rightarrow G$ exists and is smooth.

We also write

$$\begin{aligned} \text{evol}_G^l(X) &= \text{evol}_G(X) := h(1), \\ \text{Evol}_G^l(X)(t) &:= \text{evol}_G^l(s \mapsto tX(ts)) = h(t), \end{aligned}$$

if h is the (unique) solution of

$$(2) \quad \begin{cases} h(0) = e \\ \partial_t h(t) = T_e(\mu_{h(t)})(X(t)) = L_{X(t)}(h(t)), \\ \text{or } \kappa^l(\partial_t h(t)) = \delta^l h(\partial_t) = X(t). \end{cases}$$

Clearly $\text{evol}^l : C^\infty(\mathbb{R}, \mathfrak{g}) \rightarrow G$ exists and is also smooth if evol^r does, since we have $\text{evol}^l(X) = \text{evol}^r(-X)^{-1}$ by 2.4.

Let us collect some easily seen properties of the evolution mappings. If $f \in C^\infty(\mathbb{R}, \mathbb{R})$, then we have

$$\begin{aligned} \text{Evol}^r(X)(f(t)) &= \text{Evol}^r(f' \cdot (X \circ f))(t) \cdot \text{Evol}^r(X)(f(0)), \\ \text{Evol}^l(X)(f(t)) &= \text{Evol}^l(X)(f(0)) \cdot \text{Evol}^l(f' \cdot (X \circ f))(t). \end{aligned}$$

If $\varphi : G \rightarrow H$ is a smooth homomorphism between regular Lie groups then the diagram

$$\begin{array}{ccc} C^\infty(\mathbb{R}, \mathfrak{g}) & \xrightarrow{\varphi'_*} & C^\infty(\mathbb{R}, \mathfrak{h}) \\ \text{evol}_G \downarrow & & \downarrow \text{evol}_H \\ G & \xrightarrow{\varphi} & H \end{array}$$

commutes, since $\partial_t \varphi(g(t)) = T\varphi \cdot T(\mu^{g(t)}) \cdot X(t) = T(\mu^{\varphi(g(t))}) \cdot \varphi' \cdot X(t)$.

Note that each regular Lie group admits an exponential mapping, namely the restriction of evol^r to the constant curves $\mathbb{R} \rightarrow \mathfrak{g}$. A Lie group is regular if and only if its universal covering group is regular.

No counter-example to the following statement is known: *All known Lie groups modelled on convenient vector spaces are regular.* Any Banach Lie group is regular since we may consider the time dependent right invariant vector field $R_{X(t)}$ on G and its integral curve $g(t)$ starting at e , which exists and depends smoothly on (a further parameter in) X . For diffeomorphism groups the evolution operator is just integration of time dependent vector fields with compact support.

3. THE GENERAL EVOLUTION OPERATOR ALIAS THE CARTAN DEVELOPMENT

3.1. The space of flat differential forms. Let G be a regular Lie group with Lie algebra \mathfrak{g} . Let (M, x_0) be a simply connected pointed possibly infinite dimensional manifold. We consider the space

$$\Omega_{\text{flat}}^1(M, \mathfrak{g}) = \{\xi \in \Omega^1(M, \mathfrak{g}) : d\xi - \frac{1}{2}[\xi, \xi]_{\wedge}^{\mathfrak{g}} = 0\}$$

of \mathfrak{g} -valued 1-forms on M which are *flat* in the sense that they obey the left Maurer-Cartan equation.

The smooth structure on the space of flat differential forms. $\Omega_{\text{flat}}^1(M, \mathfrak{g})$ inherits the smooth structure of a Frölicher space (see [13, Section 23]) from the convenient vector space $\Omega^1(M, \mathfrak{g})$, generated by all curves $c : \mathbb{R} \rightarrow \Omega_{\text{flat}}^1(M, \mathfrak{g})$ which are smooth into $\Omega^1(M, \mathfrak{g})$. This allows us to talk about smooth mappings running through it.

Moreover, it is a Frölicher space with further structure: its kinematic tangent spaces are given by $T_{\xi}\Omega^1(M, \mathfrak{g}) = \{\eta \in \Omega^1(M, \mathfrak{g}) : d\eta - [\xi, \eta]_{\wedge}^{\mathfrak{g}} = 0\}$, and the tangent bundle is again the Frölicher space

$$T\Omega^1(M, \mathfrak{g}) = \{(\xi, \eta) \in \Omega^1(M, \mathfrak{g})^2 : d\xi - \frac{1}{2}[\xi, \xi]_{\wedge}^{\mathfrak{g}} = 0 \text{ and } d\eta - [\xi, \eta]_{\wedge}^{\mathfrak{g}} = 0\}.$$

The space $\Omega_{\text{flat}}^1(M, \mathfrak{g})$ is a manifold in the sense of [16, 17] where a cartesian closed category of manifolds based on smooth curves instead of charts as developed which up to those whose tangent spaces (proved only for finite dimensional spaces there - for Banach spaces the same proof applies) consists exactly of the usual ones.

Question: If M is finite dimensional, is $\Omega_{\text{flat}}^1(M, \mathfrak{g})$ a (split) submanifold of $\Omega^1(M, \mathfrak{g})$? Can this be shown by a quasilinear version of Hodge theory?

But we shall see below that $\Omega_{\text{flat}}^1(M, \mathfrak{g})$ is diffeomorphic to $C^\infty((M, x_0), (G, e))$ and that it gets a Lie group structure via this map, at least in the case when M is compact.

3.2. The space $C^\infty(M, G)$. If M is a compact manifold, then $C^\infty(M, G)$ is a smooth manifold; see [13, Section 42]. The same is true if M is locally compact for the space $C_c^\infty(M, G)$ of smooth mappings which equal the constant e off some compact subset of M ; let us call these smooth mappings with compact support. But in general, there is no smooth structure admitting an atlas on $C^\infty(M, G)$, since the space $C^\infty(M, N)$ is not locally contractible in its natural topology. Thus we consider $C^\infty(M, G)$ as a Frölicher Lie group with pointwise group operations in the general situation; see [13, Section 23]: The canonical smooth structure is described by

$$C^\infty(M, G) \xrightarrow{C^\infty(c, f)} C^\infty(\mathbb{R}, \mathbb{R}) \xrightarrow{\lambda} \mathbb{R}$$

where $c \in C^\infty(\mathbb{R}, M)$, where f is in $C^\infty(G, \mathbb{R})$ or in a generating set of functions, and where $\lambda \in C^\infty(\mathbb{R}, \mathbb{R})'$. It is a Frölicher space with a natural tangent bundle

$$TC^\infty(M, G) = C^\infty(M, TG) \xrightarrow{(\pi_G, \kappa^r)_*} C^\infty(M, G) \times C^\infty(M, \mathfrak{g}).$$

The space $C^\infty(M, G)$ is a manifold based on smooth curves instead of charts as developed in [16, 17].

3.3. Theorem. *Let G be a regular Lie group with Lie algebra \mathfrak{g} . Let (M, x_0) be a simply connected pointed possibly infinite dimensional manifold. In general, there exists a unique smooth mapping, called evolution operator,*

$$\text{Evol} = \text{Evol}_G^M : \Omega_{\text{flat}}^1(M, \mathfrak{g}) \rightarrow C^\infty(M, G)$$

which satisfies

$$\delta^r \circ \text{Evol} = \text{Id} \text{ and } \text{Evol}(x_0) = e.$$

Evol is a natural transformation $\Omega_{\text{flat}}^1(\cdot, \text{Lie}(\cdot)) \rightarrow C^\infty(\cdot, \cdot)$ between the two contra-covariant functors from the category of simply connected smooth manifolds times the category of Lie groups into the category of Frölicher spaces: For smooth $h : (N, y_0) \rightarrow (M, x_0)$ and $\varphi : G \rightarrow H$ a smooth homomorphism between regular Lie groups with $T_e\varphi = \varphi' : \mathfrak{g} \rightarrow \mathfrak{h}$ we have

$$C^\infty(h, \varphi) \circ \text{Evol}_G^{(M, x_0)} = \text{Evol}_H^{(N, y_0)} \circ \Omega_{\text{flat}}^1(h, \varphi');$$

in detail $\text{Evol}(h^\xi) = \text{Evol}(\xi) \circ h$ and $\varphi \circ \text{Evol}_G(\xi) = \text{Evol}_H(\varphi' \circ \xi)$.*

If one wants to avoid choosing a point x_0 in M , then Evol exists and is unique up to right translations in G . Note that a Lie group G is regular if the theorem holds for $(M, x_0) = (\mathbb{R}, 0)$. See [13, 40.2] for the main part of the proof. For finite dimensional M and Lie groups a proof can be found in [26, 27, 28], or in [2, 3] or [8] (proved with moving frames); see also [1, 5.2].

Proof. See [13, 40.2 using 39.1 and 39.2] with a small gap for the first part. For completeness' sake we repeat the proof in the more simple situation here.

If we are given a 1-form $\xi \in \Omega^1(M, \mathfrak{g})$ with $d\xi - \frac{1}{2}[\xi, \xi]_\lambda^\mathfrak{g} = 0$ then we consider the 1-form $\omega^\xi \in \Omega^1(M \times G, \mathfrak{g})$ given by

$$\omega^\xi = \kappa^l - (\text{Ad} \circ \nu) \cdot \xi, \quad \omega_{(x, g)}^\xi(Y_x, T\mu_g \cdot X) = X - \text{Ad}(g^{-1}) \cdot \xi_x(Y_x)$$

for $Y_x \in T_x M$ and $X \in \mathfrak{g}$. Then ω^ξ is a principal connection form on $M \times G$, since it reproduces the generators in \mathfrak{g} of the fundamental vector fields for the principal right action, i.e., the left invariant vector fields $0 \times L_X$, and ω^ξ is G -equivariant:

$$\begin{aligned} ((\mu^g)^* \omega^\xi)_{(x, h)} &= \omega_{(x, hg)}^\xi \circ (\text{Id}_{TM} \times T(\mu^g)) = T(\mu_{g^{-1}h^{-1}}) \cdot T(\mu^g) - \text{Ad}(g^{-1}h^{-1}) \cdot \xi_x \\ &= \text{Ad}(g^{-1}) \cdot (\kappa_h^l - \text{Ad}(h^{-1}) \cdot \xi_x) = \text{Ad}(g^{-1}) \cdot \omega_{(x, h)}^\xi. \end{aligned}$$

Since the structure group G is regular, for each smooth curve $c : \mathbb{R} \rightarrow M$, the smooth lift of c given by

$$t \mapsto (c(t), \text{Evol}_G^r(-\omega^\xi(c', 0))(t) \cdot u) = (c(t), g(t)) =: \text{Pt}^\xi(c, t, u)$$

defines a mapping $\text{Pt}^\xi(c, t, \cdot) : \{c(0)\} \times G \rightarrow \{c(t)\} \times G$ which has all the usual properties of parallel transport, namely: (1) It is horizontal since

$$\begin{aligned} g'(t) &= \partial_t \text{Evol}_G^r(-\omega^\xi(c', 0))(t) \cdot u = -T\mu^{g(t)} \cdot \omega^\xi(c', 0) \quad \text{by 2.5 (1),} \\ &= 0 + T\mu_{g(t)} \text{Ad}(g(t)^{-1}) \cdot \xi(c'(t)) \end{aligned}$$

$$\omega^\xi(c'(t), g'(t)) = \text{Ad}(g(t)^{-1}) \cdot \xi(c'(t)) - \text{Ad}(g(t)^{-1}) \cdot \xi(c'(t)) = 0.$$

(2) It is G -equivariant for the principal right action. (3) For smooth $f : \mathbb{R} \rightarrow \mathbb{R}$ we have $\text{Pt}^\xi(c, f(t), u) = \text{Pt}^\xi(c \circ f, t, \text{Pt}^\xi(c, f(0), u))$. (4) It is smooth as a mapping $\text{Pt}^\xi : C^\infty(\mathbb{R}, M) \times_{\text{ev}_0, M, \text{pr}_3} (\mathbb{R} \times M \times G) \rightarrow M \times G$ where $C^\infty(\mathbb{R}, M)$ carries the

Frölicher structure, corresponding to the compact C^∞ -topology. (5) Pt^ξ depends also smoothly on the choice of $\xi \in \Omega_{\text{flat}}^1(M, \mathfrak{g})$; this was not checked in [13, 39.1]. A smooth curve $s \mapsto \xi(s) \in \Omega_{\text{flat}}^1(M, \mathfrak{g})$ leads to a smooth curve $s \mapsto \text{Pt}^{\xi(s)}(c, t, u) \in M \times G$ by looking at the explicit formulas above.

The connection ω^ξ is flat since

$$\begin{aligned} d\omega^\xi + \frac{1}{2}[\omega^\xi, \omega^\xi]_\wedge &= d\kappa^l + \frac{1}{2}[\kappa^l, \kappa^l]_\wedge - d(\text{Ad} \circ \nu) \wedge \xi - (\text{Ad} \circ \nu).d\xi \\ &\quad - [\kappa^l, (\text{Ad} \circ \nu).\xi]_\wedge + \frac{1}{2}[(\text{Ad} \circ \nu).\xi, (\text{Ad} \circ \nu).\xi]_\wedge \\ &= -(\text{Ad} \circ \nu).(d\xi - \frac{1}{2}[\xi, \xi]_\wedge) = 0. \end{aligned}$$

Since the structure group G is regular, by theorem [13, 39.2] the horizontal bundle $\ker(\omega^\xi) \subset T(M \times G)$ is integrable: For each $x \in M$ let $u : U \rightarrow E$ be a chart with $x \in U$ and $u(x) = 0$ such that $u(U) \subset E$ is a star-shaped c^∞ -open set in E . For $y \in U$ let $c_y : [0, 1] \rightarrow U$ be given by $c_y(t) = u^{-1}(t, u(y))$. Then $\psi : U \rightarrow M \times G$ given by $\psi(y) = \text{Pt}^\xi(c_y, 1, g)$ is a smooth section $U \rightarrow M \times G$ through (x, g) which is horizontal thus an integral submanifold of $\ker \omega^\xi$; for the quite lengthy proof of horizontality see the proof of theorem [13, 39.2].

The projection $\text{pr}_1 : M \times G \rightarrow M$, restricted to each horizontal leaf, is a covering. Thus, it may be inverted over the simply connected manifold M , and the inverse $(\text{Id}_M, f) : M \rightarrow M \times G$ is a horizontal section, i.e., $T(\text{Id}_M, f) : TM \rightarrow \ker(\omega^\xi)$ is an isomorphism. Therefore

$$\begin{aligned} 0 &= ((\text{Id}, f)^*\omega^\xi)_x = (f^*\kappa^l - f^*(\text{Ad} \circ \nu).\xi)_x = (\delta^l f)_x - \text{Ad}(f(x)^{-1}).\xi_x \\ &(\delta^r f)_x = \text{Ad}(f(x)).(\delta^l f)_x = \xi_x \end{aligned}$$

for $x \in M$, as required. Moreover, $(\text{Id}_M \times f)$ is unique up to the choice of the branch of the covering and the choice of the leaf, i.e., f is unique up to a right translation by an element of G . We may fix $f =: \text{Evol}_G^M(\xi)$ by stipulating $f(x_0) = e$. Moreover, $\text{Evol}_G^M(\xi)$ depends smoothly on ξ since the parallel transport Pt^ξ depends smoothly on ξ .

It remains to check that $((M, x_0), G) \mapsto \text{Evol}_G^{(M, x_0)}$ is a natural transformation. For $h : (N, y_0) \rightarrow (M, x_0)$ we have, using 2.3

$$\delta^r \text{Evol}_G^{(N, y_0)}(h^*\xi) = h^*\xi = h^*\delta^r \text{Evol}_G^{(M, x_0)}(\xi) = \delta^r(\text{Evol}_G^{(M, x_0)}(\xi) \circ f),$$

thus $\text{Evol}_G^{(N, y_0)}(h^*\xi) = \text{Evol}_G^{(M, x_0)}(\xi) \circ f$ since both sides agree on y_0 . If $\varphi : G \rightarrow H$ is a smooth homomorphism, then $\varphi^*\kappa_H^r = \varphi'.\kappa_G^r$ since

$$\begin{aligned} (\varphi^*\kappa_H^r)_g(T_e\mu^g.X) &= (\kappa_H^r)_{\varphi(g)}(T_g\varphi.T_e\mu^g.X) = T\mu^{\varphi(g)^{-1}}.T_g\varphi.T_e\mu^g.X \\ &= T_e\varphi.X = \varphi'.T_g\mu^g.X = \varphi'.(\kappa_G^r)_g(T_e\mu^g.X), \end{aligned}$$

and obviously $\varphi'_* : \Omega_{\text{flat}}^1(M, \mathfrak{g}) \rightarrow \Omega_{\text{flat}}^1(M, \mathfrak{h})$, thus by 2.3 again we get

$$\delta^r(\varphi \circ \text{Evol}_G^M(\xi)) = \varphi'.\delta^r \text{Evol}_G^M(\xi) = \varphi'.\xi = \delta^r \text{Evol}_H^M(\varphi'.\xi)$$

thus $\varphi \circ \text{Evol}_G^M(\xi) = \text{Evol}_H^M(\varphi'.\xi)$ since both sides map $x_0 \in M$ to $e \in H$. \square

3.4. Theorem. For a regular Lie group G and simply connected (M, x_0) we have for $\xi, \eta \in \Omega_{\text{flat}}^1(M, \mathfrak{g})$:

$$\text{Evol}(\xi) \cdot \text{Evol}(\eta) = \text{Evol}\left(\xi + (\text{Ad}_G \circ \text{Evol}(\xi)) \cdot \eta\right),$$

$$\text{Evol}(\xi)^{-1} = \text{Evol}\left(-\text{Ad}_G \circ \nu \circ \text{Evol}(\xi)\right) \cdot \xi,$$

so that $\text{Evol} : \Omega_{\text{flat}}^1(M, \mathfrak{g}) \rightarrow C^\infty((M, x_0), (G, e))$ is a bijective smooth homomorphism of Frölicher Lie groups, where $C^\infty((M, x_0), (G, e))$ carries the pointwise group operations and where on $\Omega_{\text{flat}}^1(M, \mathfrak{g})$ the operations are given by

$$(\xi * \eta)(x) = \xi(x) + \text{Ad}_G(\text{Evol}(\xi)(x)) \cdot \eta(x),$$

$$\xi^{-1}(x) = -\text{Ad}_G(\text{Evol}(\xi)(x)^{-1}) \cdot \xi(x).$$

With these operations and with 0 as unit element $(\Omega_{\text{flat}}^1(M, \mathfrak{g}), *)$ becomes a regular Frölicher Lie group isomorphic to $C^\infty((M, x_0), (G, e))$. Its Lie algebra is $T_0\Omega_{\text{flat}}^1(M, \mathfrak{g}) = \{\eta \in \Omega^1(M, \mathfrak{g}) : d\eta = 0\} =: Z(M, \mathfrak{g})$ with bracket

$$[\xi_1, \xi_2]^{Z(M, \mathfrak{g})} = [\xi_1, d^{-1}\xi_2]^{\mathfrak{g}} + [d^{-1}\xi_1, \xi_2]^{\mathfrak{g}}, \quad \xi_i \in Z(M, \mathfrak{g}),$$

where

$$d^{-1} : \{\xi \in \Omega^1(M, \mathfrak{g}) : d\xi = 0\} =: Z(M, \mathfrak{g}) \rightarrow C^\infty((M, x_0), (\mathfrak{g}, 0))$$

is the bounded operator of the Poincaré lemma, the inverse of the exterior derivative.

This formula for the Lie bracket fits nicely to the 1-dimensional version derived in [13, 38.12].

Proof. We have $\mu^* \kappa^r = \text{pr}_1^* \kappa^r + (\text{Ad} \circ \text{pr}_1) \text{pr}_2^* \kappa^r$ because

$$\begin{aligned} (\mu^* \kappa^r)_{(a,b)}(\xi_a, \eta_b) &= \kappa_{\mu(ab)}^r(T\mu \cdot (\xi_a, \eta_b)) = \kappa_{ab}^r(T_a \mu^b \cdot \xi_a + T_b \mu_a \cdot \eta_b) \\ &= T_{ab} \mu^{(ab)^{-1}}(T_a \mu^b \cdot \xi_a + T_b \mu_a \cdot \eta_b) \\ &= T \mu^{a^{-1}} \cdot T \mu^{b^{-1}} \cdot T \mu^b \cdot \xi_a + T \mu^{a^{-1}} \cdot T \mu^{b^{-1}} \cdot T \mu_a \cdot \eta_b \\ &= \kappa_a^r(\xi_a) + \text{Ad}(a) \cdot \kappa_b^r(\eta_b). \end{aligned}$$

For $\xi, \eta \in \Omega_{\text{flat}}^1(M, \mathfrak{g})$ we have therefore

$$\begin{aligned} \delta^r(\text{Evol}(\xi) \cdot \text{Evol}(\eta)) &= (\mu \circ (\text{Evol}(\xi), \text{Evol}(\eta)))^* \kappa^r \\ &= (\text{Evol}(\xi), \text{Evol}(\eta))^* \mu^* \kappa^r = (\text{Evol}(\xi), \text{Evol}(\eta))^* (\text{pr}_1^* \kappa^r + (\text{Ad} \circ \text{pr}_1) \text{pr}_2^* \kappa^r) \\ &= \text{Evol}(\xi)^* \kappa^r + (\text{Ad} \circ \text{Evol}(\xi)) \cdot \text{Evol}(\eta)^* \kappa^r = \xi + \text{Ad}(\text{Evol}(\xi)) \eta. \end{aligned}$$

which implies

$$\text{Evol}(\xi) \cdot \text{Evol}(\eta) = \text{Evol}(\xi * \eta), \quad \text{Evol}(\xi)^{-1} = \text{Evol}(\xi^{-1}).$$

Thus, $\text{Evol} : \Omega_{\text{flat}}^1(M, \mathfrak{g}) \rightarrow C^\infty(M, G)$ is a group isomorphism onto the subgroup $C^\infty((M, x_0), (G, e)) := \{f \in C^\infty(M, G) : f(x_0) = e\}$ with the pointwise product, which, however, is only a Frölicher space in general, see [13, 23.1]. If M is compact then $C^\infty((M, x_0), (G, e))$ is smooth regular Lie group. If M is finite dimensional, then one has to refine the topology (control near infinity) to make it into a disjoint union of regular Lie groups, and then one has to mimick this procedure also on $\Omega_{\text{flat}}^1(M, \mathfrak{g})$.

In general, we will just take both groups as Frölicher spaces with special properties (having a tangent bundle, e.g.).

It follows that the product on $\Omega_{\text{flat}}^1(M, \mathfrak{g})$ has the properties of a group structure.

A direct proof is fun and needs naturality of Evol in an essential way. As entertainment we compute the following:

$$\begin{aligned} \xi^{-1} * \xi &= -\text{Ad}_G(\text{Evol}(\xi)^{-1}).\xi + \text{Ad}_G(\text{Evol}(-\text{Ad}_G(\text{Evol}(\xi)^{-1}).\xi).\xi) = 0 \text{ since} \\ &(\text{Evol}(\xi)^{-1})^* \kappa^r = (\nu \circ \text{Evol}(\xi))^* \kappa^r = \text{Evol}(\xi)^* \nu^* \kappa^r \\ &= \text{Evol}(\xi)^*(-\text{Ad} \circ \nu) \kappa^r = -\text{Ad}(\text{Evol}(\xi)^{-1}) \text{Evol}(\xi)^* \kappa^r \\ &= -\text{Ad}(\text{Evol}(\xi)^{-1})\xi \text{ implies again the expression for the inverse} \\ &\text{Evol}(\xi)^{-1} = \text{Evol}(-\text{Ad}(\text{Evol}(\xi)^{-1})\xi). \end{aligned}$$

Now we aim for the Lie bracket. Since $\delta^r : C^\infty((M, x_0), (G, e)) \rightarrow \Omega_{\text{flat}}^1(M, \mathfrak{g}) \subset \Omega^1(M, \mathfrak{g})$ is a smooth group isomorphism of Frölicher Lie groups, we can just take its tangent mapping at the constant e which will become a homomorphism of Lie algebras. As an aside note that δ^r is a Lie derivative analogous to the finite dimensional version in [12, Chapter IX]. The infinite dimensional version has been worked out in detail in [15, 12.2–12.5] for mappings $f : M \rightarrow N$; here the only difference is that the forms are \mathfrak{g} -valued and that we make use the right trivialization of TG . So we choose a smooth curve $t \mapsto f(t) \in C^\infty((M, x_0), (G, e))$ with $f(0) = e$ and $\partial_t|_0 f(t) = X \in C^\infty(M, \mathfrak{g})$ with $X(x_0) = 0$; then we consider the smooth mapping $\hat{f} : \mathbb{R} \times M \rightarrow G$. By the Maurer-Cartan equation 2.3, for a vector field $Y \in \mathfrak{X}(M)$ we have

$$\begin{aligned} 0 &= d(\delta^r \hat{f})(\partial_t, 0_M), (0_{\mathbb{R}}, Y) - [(\delta^r \hat{f})(\partial_t, 0_M), (\delta^r \hat{f})(0_{\mathbb{R}}, Y)]^{\mathfrak{g}} \\ &= \partial_t((\delta^r \hat{f})(0_{\mathbb{R}}, Y)) - \mathcal{L}_Y(T\mu^{f^{-1}} \partial_t f) - (\delta^r \hat{f})([(\partial_t, 0_M), (0_{\mathbb{R}}, Y)]^{\mathfrak{X}(\mathbb{R} \times M)}) \\ &\quad - [(\delta^r \hat{f})(\partial_t, 0_M), (\delta^r \hat{f})(0_{\mathbb{R}}, Y)]. \end{aligned}$$

Choosing $t = 0$ and using $f(0, x) = e$ this becomes

$$(T_e \delta^r . X)(Y) = \mathcal{L}_Y X = dX(Y).$$

Now we can write down the Lie bracket. Since M is simply connected, its de Rham cohomology $H^1(M) = 0$, which also holds for \mathfrak{g} -valued 1-forms and for infinite dimensional M ; see [13, Section 34]. Let

$$d^{-1} : \{\xi \in \Omega^1(M, \mathfrak{g}) : d\xi = 0\} =: Z(M, \mathfrak{g}) \rightarrow C^\infty((M, x_0), (\mathfrak{g}, 0))$$

be the bounded (by lemma 3.5 below) operator of the Poincaré lemma; i.e., the inverse of exterior derivative. Then for $X_i \in C^\infty((M, x_0), (\mathfrak{g}, 0))$ we have

$$\begin{aligned} T_e \delta^r . [X_1, X_2]^{\mathfrak{g}} &= d[X_1, X_2]^{\mathfrak{g}} = [dX_1, X_2]^{\mathfrak{g}} + [X_1, dX_2]^{\mathfrak{g}}, \quad \text{thus} \\ [X_1, X_2]^{\mathfrak{g}} &= [d^{-1} \xi_1, d^{-1} \xi_2]^{\mathfrak{g}} + [d^{-1} \xi_1, \xi_2]^{\mathfrak{g}}, \quad \text{for } \xi_i \in Z(M, \mathfrak{g}). \quad \square \end{aligned}$$

3.5. Lemma. *For a simply connected manifold M the inverse of the exterior derivative $d^{-1} : Z(M, \mathfrak{g}) \rightarrow C^\infty((M, x_0), (\mathfrak{g}, 0))$ is a bounded linear operator.*

Proof. For a star-shaped c^∞ -open subset M in a convenient vector space E this follows from the explicit formula for the Poincaré operator $d^{-1}\xi(x) = \int_0^1 t\omega(tx)(x)dt$.

In general, we cover the simply connected manifold M by open charts $u : U \rightarrow u(U) \subset E$ with $x_0 \in U$, $u(x_0) = 0$, and $u(U)$ star-shaped in E . Via the diffeomorphisms u^* the operators $d^{-1} : Z(U, \mathfrak{g}) \rightarrow C^\infty((U, x_0), (\mathfrak{g}, 0))$ are all bounded. Since $C^\infty((M, x_0), (\mathfrak{g}, 0))$ and $Z(M, \mathfrak{g})$ both carry the initial structure with respect to the restriction operators to U , the result follows. \square

3.6. Theorem. *For a regular Lie group and $\xi \in \Omega_{flat}^1(M, \mathfrak{g})$ we consider $f = \text{Evol}(\xi) : (M, x_0) \rightarrow (G, e)$. For a mapping $h : M \rightarrow \mathfrak{g}$ we consider $\eta := \text{Ad}(f)dh \in \Omega^1(M, \mathfrak{g})$. Then we have $d\eta - \frac{1}{2}[\xi, \eta]_\wedge^\mathfrak{g} = 0$.*

If conversely $\eta \in \Omega^1(M, \mathfrak{g})$ satisfies $d\eta - \frac{1}{2}[\xi, \eta]_\wedge^\mathfrak{g} = 0$ then there exists a unique smooth $h : (M, x_0) \rightarrow (\mathfrak{g}, 0)$ such that $dh = \text{Ad}(f^{-1})\eta$.

Proof. If $h : M \rightarrow \mathfrak{g}$ exists then $d^2h = 0$ so that

$$\begin{aligned} d\eta &= d\text{Ad}.Tf \wedge dh + \text{Ad}(f)d^2h = (\text{ad} \circ \kappa^r.Tf) \text{Ad}(f)dh + 0 \\ &= \frac{1}{2}[f^*\kappa^r, \eta]_\wedge^\mathfrak{g} = \frac{1}{2}[\xi, \eta]_\wedge^\mathfrak{g}. \end{aligned}$$

Conversely, by this computation $d\eta - \frac{1}{2}[\xi, \eta]_\wedge^\mathfrak{g} = 0$ implies that the \mathfrak{g} -valued 1-form $\beta := \text{Ad}(f^{-1})\eta$ is closed. Since M is simply connected, $\beta = dh$ for $h \in C^\infty(M, \mathfrak{g})$ which is unique up to addition of a constant. \square

3.7. Theorem. *Let G be a Lie group. Then via right trivialization $(\kappa^r, \pi_G) : TG \rightarrow \mathfrak{g} \times G$ the tangent group TG is isomorphic to the semidirect product $\mathfrak{g} \rtimes G$, where G acts by $\text{Ad} : G \rightarrow \text{Aut}(\mathfrak{g})$. So for $g, h \in G$ and $X, Y \in \mathfrak{g}$ we have:*

$$\begin{aligned} (1) \quad \mu_{\mathfrak{g} \rtimes G}((X, g), (Y, h)) &= (X + \text{Ad}(g)Y, gh) \\ \nu_{\mathfrak{g} \rtimes G}(X, g) &= (-\text{Ad}(g^{-1})X, g^{-1}), \\ [(X_1, Y_1), (X_2, Y_2)]_{\mathfrak{g} \rtimes \mathfrak{g}} &= ([Y_1, X_2] - [Y_2, X_1], [Y_1, Y_2]), \\ \text{Ad}_{(X, g)}^{\mathfrak{g} \rtimes G}(Y, Z) &= (\text{Ad}(g)Y - [\text{Ad}(g)Z, X], \text{Ad}(g)Z) \end{aligned}$$

If G is a regular Lie group, then so is $TG \cong \mathfrak{g} \rtimes G$ and $T\text{evol}_G^{\mathbb{R}}$ corresponds to $\text{evol}_{TG}^{\mathbb{R}}$, via

$$(2) \quad \begin{array}{ccc} TC^\infty(\mathbb{R}, \mathfrak{g}) & \xrightarrow{\cong} & C^\infty(\mathbb{R}, \mathfrak{g} \rtimes \mathfrak{g}) \\ T\text{evol}_G^{\mathbb{R}} \downarrow & \swarrow \text{evol}_{TG}^{\mathbb{R}} & \downarrow \text{evol}_{\mathfrak{g} \rtimes G}^{\mathbb{R}} \\ TG & \xrightarrow{\cong} & \mathfrak{g} \rtimes G. \end{array}$$

In particular, for $(Y, X) \in C^\infty(\mathbb{R}, \mathfrak{g} \rtimes \mathfrak{g}) = TC^\infty(\mathbb{R}, \mathfrak{g})$, where X is the footpoint, we have

$$\begin{aligned} \text{evol}_{\mathfrak{g} \rtimes G}^{\mathbb{R}}(Y, X) &= \left(\text{Ad}(\text{evol}_G^{\mathbb{R}}(X)) \int_0^1 \text{Ad}(\text{Evol}_G^{\mathbb{R}}(X)(s)^{-1}).Y(s) ds, \text{evol}_G^{\mathbb{R}}(X) \right) \\ (3) \quad T_X \text{evol}_G^{\mathbb{R}}.Y &= T(\mu_{\text{evol}_G^{\mathbb{R}}(X)}). \int_0^1 \text{Ad}(\text{Evol}_G^{\mathbb{R}}(X)(s)^{-1}).Y(s) ds, \\ T_X(\text{Evol}_G^{\mathbb{R}}(\quad)(t)).Y &= T(\mu_{\text{Evol}_G^{\mathbb{R}}(X)(t)}). \int_0^t \text{Ad}(\text{Evol}_G^{\mathbb{R}}(X)(s)^{-1}).Y(s) ds. \end{aligned}$$

For a pointed simply connected manifold (M, x_0) we have

$$(4) \quad \begin{aligned} T\Omega_{\text{flat}}^1(M, \mathfrak{g}) &= \{(\xi, \eta) \in \Omega^1(M, \mathfrak{g})^2 : d\xi - \frac{1}{2}[\xi, \xi]_{\wedge}^{\mathfrak{g}} = 0 \text{ and } d\eta - [\xi, \eta]_{\wedge}^{\mathfrak{g}} = 0\} \\ &\cong \{(\eta, \xi) \in \Omega^1(M, \mathfrak{g} \times \mathfrak{g}) : (d\eta - [\xi, \eta]_{\wedge}^{\mathfrak{g}}, d\xi - \frac{1}{2}[\xi, \xi]_{\wedge}^{\mathfrak{g}}) = (0, 0)\} \\ &= \Omega_{\text{flat}}^1(M, \mathfrak{g} \times \mathfrak{g}) \end{aligned}$$

and the following diagram commutes:

$$(5) \quad \begin{array}{ccc} T\Omega_{\text{flat}}^1(M, \mathfrak{g}) & \xrightarrow{T \text{Evol}_G^M} & TC^\infty((M, x_0), (G, e)) \\ \cong \downarrow & & \downarrow \cong \\ \Omega_{\text{flat}}^1(M, T\mathfrak{g}) & \xrightarrow{\text{Evol}_{TG}^M} & C^\infty((M, x_0), (TG, 0_e)) \\ \cong \downarrow & & \downarrow \cong \\ \Omega_{\text{flat}}^1(\mathbb{R}, \mathfrak{g} \times \mathfrak{g}) & \xrightarrow{\text{Evol}_{\mathfrak{g} \times G}^M} & C^\infty((M, x_0), (\mathfrak{g} \times G, (0, e))). \end{array}$$

Proof. The first part may be found in [13, 38.10]. We repeat the derivation of the formulas. By 2.1, for $g, h \in G$ and $X, Y \in \mathfrak{g}$, we have

$$\begin{aligned} T_{(g,h)}\mu.(R_X(g), R_Y(h)) &= T(\mu^h).R_X(g) + T(\mu_g).R_Y(h) \\ &= T(\mu^h).T(\mu^g).X + T(\mu_g).T(\mu^h).Y = R_X(gh) + R_{\text{Ad}(g)Y}(gh), \\ T_g\nu.R_X(g) &= -T(\mu^{g^{-1}}).T(\mu_{g^{-1}}).T(\mu^g).X = -R_{\text{Ad}(g^{-1})X}(g^{-1}), \end{aligned}$$

so that μ_{TG} and ν_{TG} , and the Lie bracket, after right trivialization, are given by

$$(6) \quad \begin{aligned} \mu^{\mathfrak{g} \times G}((X, g), (Y, h)) &= (X + \text{Ad}(g)Y, gh) \\ T\mu_{(X,g)}^{\mathfrak{g} \times G}(Y', h') &= (\text{Ad}(g)Y', T\mu_g h'), \quad h' \in T_h G \\ \nu^{\mathfrak{g} \times G}(X, g) &= (-\text{Ad}(g^{-1})X, g^{-1}), \\ (X, g).(Y, h).(X, g)^{-1} &= (X + \text{Ad}(g)Y, gh).(-\text{Ad}(g^{-1})X, g^{-1}) \\ &= (X + \text{Ad}(g)Y - \text{Ad}(ghg^{-1})X, ghg^{-1}) \\ \text{Ad}^{\mathfrak{g} \times G}(X, g)(Y', h') &= (\text{Ad}(g)Y' - \text{Ad}(e)[\text{Ad}(g)h', X], \text{Ad}(g)h') \\ \text{ad}^{\mathfrak{g} \times G}(X', g')(Y', h') &= ([g', Y'] - [[g', h'], 0]) - [h', X'], [g', h'] \\ [(X_1, Y_1), (X_2, Y_2)]_{\mathfrak{g} \times \mathfrak{g}} &= ([Y_1, X_2] - [Y_2, X_1], [Y_1, Y_2]). \end{aligned}$$

That diagram (2) commutes and equations (3) hold, has been proven in [13, 38.10]. Now we prove that diagram (5) commutes. For the bottom square this follows from theorem 3.3. We consider a curve $t \mapsto \xi(t) \in \Omega_{\text{flat}}^1(M, \mathfrak{g})$ which is smooth into $\Omega^1(M, \mathfrak{g})$. Then $\partial_t|_0 \xi(t) =: \eta \in T_{\xi(0)}\Omega_{\text{flat}}^1(M, \mathfrak{g})$ so that $d\eta - [\xi(0), \eta]_{\wedge}^{\mathfrak{g}} = 0$. As in the beginning of the proof of theorem 3.3 we now consider the smooth curve of flat principal connections

$$t \mapsto \omega^{\xi(t)} = \kappa^l - (\text{Ad} \circ \nu).\xi(t)$$

on $M \times G$, and we let $L^{\xi(t)} \subset M \times G$ be the horizontal leaf through (x_0, e) for the connection $\omega^{\xi(t)}$. Recall that

$$(\text{Id}_M, \text{Evol}_G^M(\xi(t))) = (\text{pr}_1 | L^{\xi(t)})^{-1} : M \rightarrow L^{\xi(t)} \subset M \times G$$

is the horizontal lift. Likewise, writing $\xi = \xi(0)$,

$$(\text{Id}_M, \text{Evol}_{\mathfrak{g} \rtimes G}^M(\eta, \xi) = (\text{pr}_1 | L^{(\eta, \xi)})^{-1} : M \rightarrow L^{(\eta, \xi)} \subset M \times (\mathfrak{g} \rtimes G)$$

where $L^{(\eta, \xi)}$ is the horizontal leaf through $(x_0, 0, e)$ of the principal connection

$$\omega^{(\eta, \xi)} = \kappa^L_{\mathfrak{g} \rtimes G} - (\text{Ad}^{\mathfrak{g} \rtimes G} \circ \nu^{\mathfrak{g} \rtimes G})(\eta, \xi).$$

Claim. $\partial_t|_0 L^{\xi(t)}$ is the horizontal leaf $L^{(\eta, \xi)}$ of the flat principal connection $\omega^{(\eta, \xi(0))}$ on the principal bundle $M \times (\mathfrak{g} \rtimes G) \rightarrow M$. This is sufficient to finish the proof.

$T_{(x_0, e)} L^{\xi(t)}$ consists of all $(Y, X(t)) \in T_{x_0} M \times \mathfrak{g}$ such that $\omega_{(x_0, e)}^{\xi(t)}(Y, X(t)) = X(t) - \xi(t)(Y) = 0$; since $L^{\xi(t)}$ is a horizontal leaf we may fix Y . Then

$$\partial_t|_0 \omega_{(x_0, e)}^{\xi(t)}(Y, X(t)) = X' - \eta(Y) = 0.$$

On the other hand $\omega_{(x_0, (0, e))}^{(\eta, \xi)}(Y, (Z, X)) = (Z - \eta(Y), X - \xi(Y))$ since for $(X, g) \in \mathfrak{g} \rtimes G$ and $(X', g') \in T_X \mathfrak{g} \times T_g G$ we have

$$\begin{aligned} \omega_{(x, (X, g))}^{(\eta, \xi)}(Y, (X, X'; g, g')) &= (T\mu_{(X, g)^{-1}}^{\mathfrak{g} \rtimes G})(X', g') - \text{Ad}^{\mathfrak{g} \rtimes G}((X, g)^{-1})(\eta(Y), \xi(Y)) \\ &= (T\mu_{(-\text{Ad}(g^{-1})X, g^{-1})}^{\mathfrak{g} \rtimes G})(X', g') - \text{Ad}^{\mathfrak{g} \rtimes G}(-\text{Ad}(g^{-1})X, g^{-1})(\eta(Y), \xi(Y)) \\ &= (\text{Ad}(g^{-1})X', T\mu_{g^{-1}} g') \\ &\quad - (\text{Ad}(g^{-1})\eta(Y) - [\text{Ad}(g^{-1})\xi(Y), -\text{Ad}(g^{-1})X], \text{Ad}(g^{-1})\xi(Y)) \\ &= (\text{Ad}(g^{-1})X' - \text{Ad}(g^{-1})\eta(Y) - \text{Ad}(g^{-1})[\xi(Y), X], T\mu_{g^{-1}} g' - \text{Ad}(g^{-1})\xi(Y)). \quad \square \end{aligned}$$

REFERENCES

- [1] D. V. Alekseevsky and P. W. Michor. Differential geometry of Cartan connections. *Publ. Math. Debrecen*, 47(3-4):349–375, 1995.
- [2] E. Cartan. La méthode du repère mobile, la théorie des groupes continus et les espaces généralisés. (Exposés de géométrie V.) Actual. scient. et industr. 1935, Nr. 194, 65 p., 1935.
- [3] E. Cartan. *La théorie des groupes finis et continus et la géométrie différentielle traitées par la méthode du repère mobile. Leçons professées à la Sorbonne. Rédigées par J. Leray.* Cahiers scientifiques. Fasc. 18. 1937.
- [4] T. Diez, A. Futaki, and T. Ratiu. Cartan geometry and infinite-dimensional Kempf-Ness theory, 2024. arXiv:2405.20864.
- [5] A. S. Fokas and I. M. Gelfand. Surfaces on Lie groups, on Lie algebras, and their integrability. *Comm. Math. Phys.*, 177(1):203–220, 1996. With an appendix by Juan Carlos Alvarez Paiva.
- [6] H. Glöckner. Regularity properties of infinite-dimensional Lie groups, and semiregularity, 2012. arXiv:1208.0715.
- [7] H. Glöckner. Measurable regularity properties of infinite-dimensional Lie groups, 2015. arXiv:1601.02568.
- [8] P. Griffiths. On Cartan’s method of Lie groups and moving frames as applied to uniqueness and existence questions in differential geometry. *Duke Math. J.*, 41:775–814, 1974.
- [9] M. Hanusch. Differentiability of the evolution map and Mackey continuity. *Forum Math.*, 31(5):1139–1177, 2019.
- [10] M. Hanusch. The regularity problem for Lie groups with asymptotic estimate Lie algebras. *Indag. Math. (N.S.)*, 31(1):152–176, 2020.
- [11] M. Hanusch. Regularity of Lie groups. *Comm. Anal. Geom.*, 30(1):53–152, 2022.
- [12] I. Kolář, P. W. Michor, and J. Slovák. *Natural operations in differential geometry.* Springer-Verlag, Berlin, 1993.

- [13] A. Kriegl and P. W. Michor. *The convenient setting of global analysis*, volume 53 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997. http://www.ams.org/online_bks/surv53/.
- [14] A. Kriegl and P. W. Michor. Regular infinite-dimensional Lie groups. *J. Lie Theory*, 7(1):61–99, 1997.
- [15] P. W. Michor. *Manifolds of differentiable mappings*, volume 3 of *Shiva Mathematics Series*. Shiva Publishing Ltd., Nantwich, 1980.
- [16] P. W. Michor. A convenient setting for differential geometry and global analysis. I. *Cahiers Topologie Géom. Différentielle*, 25(1):63–109, 1984.
- [17] P. W. Michor. A convenient setting for differential geometry and global analysis. II. *Cahiers Topologie Géom. Différentielle*, 25(2):113–178, 1984.
- [18] P. W. Michor. Manifolds of mappings for continuum mechanics. In *Geometric Continuum Mechanics*, volume 42 of *Advances in Continuum Mechanics*, pages 3–75. Birkhäuser Basel, 2020.
- [19] J. Milnor. Remarks on infinite-dimensional Lie groups. In *Relativity, groups and topology, II (Les Houches, 1983)*, pages 1007–1057. North-Holland, Amsterdam, 1984.
- [20] H. Omori, Y. Maeda, and A. Yoshioka. On regular Fréchet Lie groups I. Some differential geometric expressions of Fourier integral operators on a Riemannian manifold. *Tokyo J. Math.*, 3:353–390, 1980.
- [21] H. Omori, Y. Maeda, and A. Yoshioka. On regular Fréchet Lie groups II. Composition rules of Fourier Integral operators on a Riemannian manifold. *Tokyo J. Math.*, 4:221–253, 1981.
- [22] H. Omori, Y. Maeda, and A. Yoshioka. On regular Fréchet Lie groups III. *Tokyo J. Math.*, 4:255–277, 1981.
- [23] H. Omori, Y. Maeda, and A. Yoshioka. On regular Fréchet Lie groups IV. Definitions and fundamental theorems. *Tokyo J. Math.*, 5:365–398, 1982.
- [24] H. Omori, Y. Maeda, and A. Yoshioka. On regular Fréchet Lie groups V. Several basic properties. *Tokyo J. Math.*, 6:39–64, 1983.
- [25] H. Omori, Y. Maeda, A. Yoshioka, and O. Kobayashi. On regular Fréchet Lie groups VI. Infinite dimensional Lie groups which appear in general relativity. *Tokyo J. Math.*, 6:217–246, 1983.
- [26] A. L. Onishchik. On the classification of fiber spaces. *Sov. Math. Doklady*, 2:1561–1564, 1961.
- [27] A. L. Onishchik. Connections with zero curvature and the de Rham theorem. *Sov. Math. Doklady*, 5:1654–1657, 1964.
- [28] A. L. Onishchik. Some concepts and applications of non-abelian cohomology theory. *Trans. Moscow Math. Soc.*, 17:49–98, 1967.
- [29] Wikipedia: Convenient vector space. Convenient vector space — Wikipedia, the free encyclopedia, 2015. [Online; accessed 25-February-2024].

PETER W. MICHOR: FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, OSKAR-MORGENSTERN-PLATZ 1, A-1090 WIEN, AUSTRIA.

Email address: Peter.Michor@univie.ac.at

JOHANNA MICHOR: FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, OSKAR-MORGENSTERN-PLATZ 1, A-1090 WIEN, AUSTRIA.

Email address: Johanna.Michor@univie.ac.at